

Adenosine type 1 (A₁) receptors mediate protection against myocardial infarction produced by chronic, intermittent ingestion of ethanol in dogs

Franz Kehl^a, John G. Krolkowski^a, John F. LaDisa Jr.^{a,d}, Judy R. Kersten^{a,c},
David C. Warltier^{a,b,c,d}, Paul S. Pagel^{a,d,*}

^aDepartment of Anesthesiology, Medical College of Wisconsin, MEB-M4280, 8701 Watertown Plank Road, Milwaukee, WI 53226, USA

^bDepartment of Medicine (Division of Cardiovascular Diseases), Medical College of Wisconsin, MEB-M4280, 8701 Watertown Plank Road, Milwaukee, WI 53226, USA

^cDepartment of Pharmacology and Toxicology, Medical College of Wisconsin, MEB-M4280, 8701 Watertown Plank Road, Milwaukee, WI 53226, USA

^dThe Clement J. Zablocki Veterans Affairs Medical Center and the Department of Biomedical Engineering, Marquette University, Milwaukee, WI, USA

Received 8 February 2002; received in revised form 16 May 2002; accepted 19 May 2002

Abstract

Background: Chronic consumption of small amounts of ethanol protects myocardium from ischemic injury. We tested the hypothesis that adenosine type 1 (A₁) receptors mediate these beneficial effects. **Methods:** Dogs ($n=37$) were fed with ethanol (1.5 g/kg) or water mixed with dry food twice per day for 12 weeks, fasted overnight before experimentation, and instrumented for measurement of hemodynamics. Dogs received intravenous drug vehicle (50% polyethylene glycol in 0.1 N sodium hydroxide and 0.9% saline over 15 min) or the selective A₁ receptor antagonist 8-cyclopentyl-1,3-dipropylxanthine (DPCPX, 0.8 mg/kg over 15 min) and were subjected to a 60 min coronary artery occlusion followed by 3 h of reperfusion. Myocardial infarct size and transmural coronary collateral blood flow were measured with triphenyltetrazolium chloride staining and radioactive microspheres, respectively. **Results:** The area at risk (AAR) for infarction was similar between groups. Pretreatment with ethanol significantly reduced infarct size to $13\pm 2\%$ ($n=7$) of the AAR as compared to control experiments ($26\pm 2\%$; $n=7$). DPCPX abolished the protective effects of ethanol pretreatment ($30\pm 3\%$; $n=7$) but had no effect in dogs that did not receive ethanol ($25\pm 2\%$; $n=7$). No differences in transmural coronary collateral blood flow were observed between groups. **Conclusions:** The present findings indicate that chronic ingestion of small amounts of ethanol produces myocardial protection that persists after the discontinuation of ethanol. The results indicate that A₁ receptors mediate ethanol-induced preconditioning in dogs independent of alterations in systemic hemodynamics or coronary collateral blood flow.

© 2002 Elsevier Science Ireland Ltd. All rights reserved.

Keywords: Myocardial infarction; Infarct size; Prolonged coronary occlusion; Myocardial ischemia; Ethanol; Pharmacologic preconditioning

1. Introduction

Chronic consumption of small amounts of ethanol improves survival in patients after acute myocardial

infarction [1,2], but the mechanisms responsible for this beneficial effect are incompletely described. We [3,4] and others [5] have recently demonstrated that chronic, moderate ethanol consumption preserves myocardium from irreversible ischemic damage by activation of mitochondrial adenosine triphosphate-dependent potassium (K_{ATP}) channels, a process that we termed ‘chronic ethanol-induced preconditioning’

*Corresponding author. Tel.: +1-414-456-5733; fax: +1-414-456-6507.

E-mail address: pspagel@mcw.edu (P.S. Pagel).

(CEPC). Mitochondrial K_{ATP} channels [6] also mediate the protective effects of ischemic preconditioning (IPC) through an intracellular signal transduction pathway that includes adenosine type 1 (A_1) receptors [7] coupled to inhibitory guanine (G_i) nucleotide binding proteins [8] and protein kinase C (PKC) [9].

Evidence initially obtained in isolated guinea pig and rat hearts indicates that chronic exposure to small amounts of ethanol protects myocardium from subsequent ischemic injury by a signal transduction pathway similar to that implicated in IPC [10–12]. The salutary action of chronic ingestion of low doses of ethanol was attributed to activation of adenosine or alpha-adrenergic receptors [10,12] and sustained translocation of the epsilon isoform of PKC [11]. However, hemodynamics and regional myocardial perfusion were not quantified nor were plasma ethanol concentrations assessed in these previous studies [10–12]. In contrast, we [3,4] have employed a unique canine model of intermittent ethanol ingestion during the pretreatment period [13] that more closely resembles moderate ethanol consumption with meals in humans and consistently produces subintoxicating plasma ethanol concentrations [3]. Acute experimentation after ethanol pretreatment in this large animal model allows invasively-derived assessment of systemic hemodynamics and precise quantification of coronary collateral blood flow before and during prolonged coronary artery occlusion and reperfusion *in vivo*. In the present investigation, we tested the hypothesis that A_1 receptors mediate the protective effects of chronic intermittent ingestion of ethanol independent of alterations in hemodynamics or transmural myocardial perfusion.

2. Methods

All experimental procedures and protocols used in this investigation were reviewed and approved by the Animal Care and Use Committee of the Medical College of Wisconsin. All conformed to the Guiding Principles in the Care and Use of Animals of the American Physiological Society and the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health (Revised, 1996).

2.1. General preparation

Mongrel dogs weighing between 25 and 30 kg were randomly assigned to receive dry dog chow (Lab Canine Diet, Richmond, IN) mixed with ethanol (1.5 g/kg) or an equal volume of water twice a day for 12 weeks [3,4,13]. Drinking water was provided *ad libitum*. We [3] have previously shown that administration of ethanol using this method produces peak blood ethanol concentrations of 52 ± 4 mg/dl 45 min after eating. Dogs that did not consume the chow–ethanol mix or lost weight during the 12 week pretreatment period were excluded from further experimentation. Dogs were fasted overnight before and did not receive ethanol on the day of experimentation.

2.2. Implantation of instruments

Dogs were anesthetized with sodium barbital (200 mg/kg) and sodium pentobarbital (15 mg/kg) and ventilated with an air and oxygen mixture (fraction of inspired oxygen=0.25) after intubation of the trachea as previously described [3,14]. Acid–base status and arterial blood gas tensions were maintained within the normal range by adjustment of respiratory rate and tidal volume throughout the experiment. After calibration, a double pressure transducer-tipped catheter was inserted into the aorta and left ventricle (LV) through the left carotid artery to measure arterial and LV pressures. The maximal rate of increase of LV pressure ($+dP/dt_{max}$) was obtained by electronic differentiation of the LV pressure wave form. The femoral artery and vein were cannulated for the withdrawal of reference blood flow samples and fluid administration, respectively. A thoracotomy was performed in the left fifth intercostal space. A heparin-filled catheter was inserted into the left atrial appendage for administration of radioactive microspheres. A 1.0 cm segment of the left anterior descending coronary artery (LAD) was dissected immediately distal to the first diagonal branch, and a silk ligature was placed around this vessel for production of coronary artery occlusion and reperfusion. The appearance of cyanosis and regional dyskinesia of myocardium immediately distal to the ligature were used to verify the adequacy of LAD occlusion in all experiments. Hemodynamic data were continuously

monitored throughout the experiment, recorded on a polygraph, and digitized using a computer interfaced with an analog-to-digital converter.

2.3. Measurement of myocardial infarct size

At the conclusion of each experiment, the LAD was again occluded and cannulated at the occlusion site [14]. Briefly, 10 ml each of saline and patent blue dye were injected at equal pressure in the LAD and left atrium to delineate the anatomic area at risk (AAR) and the normal zone, respectively. The heart was fibrillated, removed, and sliced into serial 6 to 7 mm wide transverse sections. The unstained AAR was separated from the normal area, and the two regions were incubated for 20 min at 37 °C in 1% 2,3,5-triphenyltetrazolium chloride in 0.1 M phosphate buffer adjusted to pH=7.4. Infarcted and noninfarcted myocardium within the AAR were separated and weighed after being stored overnight in 10% formaldehyde. Infarct size was expressed as a percentage of AAR.

2.4. Measurement of regional myocardial perfusion

Carbonized plastic microspheres [15±2 µm (SD)] labeled with ⁹⁵Nb, ¹⁴¹Ce, and ¹⁰³Ru were used to measure myocardial perfusion as previously described [14]. Briefly, microspheres were administered into the left atrium as a bolus and flushed in with 10 ml of warm (37 °C) saline. A few seconds before injection, a timed collection of reference arterial blood flow was started from the femoral arterial catheter at a rate of 7 ml/min for 3 min. Microspheres were injected 5 min before LAD occlusion (after administration of vehicle or DPCPX), 30 min after LAD occlusion, and 1 h after final reperfusion. Transmural tissue samples were selected from the ischemic region and subdivided into subepicardial, midmyocardial, and subendocardial layers of approximately equal thickness. Samples were weighed and placed in scintillation vials, and the activity of each isotope was determined. Similarly, the activity of each isotope in the reference blood flow sample was assessed. Tissue blood flow (ml/min per g) was calculated as $Q_r \cdot C_m / C_r$, where Q_r is the rate of withdrawal of the reference blood flow sample (ml/

min), C_m is the activity (cpm/g) of the myocardial tissue sample, and C_r is the activity of the reference blood flow sample. Transmural blood flow was considered to be the average of the subepicardial, midmyocardial, and subendocardial blood flows.

2.5. Experimental protocol

Baseline hemodynamics were recorded 90 min after completion of the surgical preparation. All dogs were subjected to a 60 min LAD occlusion followed by 3 h of reperfusion. In four separate groups of experiments, dogs that consumed ethanol or water (control) mixed with dog chow for 12 weeks were randomly assigned to receive intravenous infusions of drug vehicle (50% polyethylene glycol in 0.1 N sodium hydroxide and 0.9% saline over 15 min) or the selective A₁ receptor antagonist 8-cyclopentyl-1,3-dipropylxanthine (DPCPX; 0.8 mg/kg over 15 min) 60 min before LAD occlusion and reperfusion. We have previously shown that this dose of DPCPX markedly attenuates enhanced functional recovery of stunned myocardium produced by the volatile anesthetic isoflurane in dogs [15]. Dogs that developed intractable ventricular fibrillation and those with a subendocardial coronary collateral blood flow=0.15 ml/min per g were excluded from subsequent data analysis [16].

2.6. Statistical analysis

Statistical analysis of data within and between groups was performed with analysis of variance (ANOVA) with repeated measures followed by Student's *t*-test with Bonferroni's correction for multiplicity. Linear regression analysis was performed to determine the relationship between transmural collateral blood flow and myocardial infarct size. Changes within and between groups were considered statistically significant when $P < 0.05$. Data are expressed as mean±standard error of the mean (S.E.M.).

3. Results

All dogs consumed the assigned diet during the 12 week pretreatment period. Thirty seven dogs were used to obtain 28 successful experiments. Four dogs

were excluded from analysis because of intractable ventricular fibrillation during LAD occlusion or reperfusion (1 control, 2 ethanol pretreatment alone, 1 ethanol pretreatment and DPCPX). Five dogs were excluded because subendocardial coronary collateral blood flow exceeded 0.15 ml/min/g (1 control, 1 DPCPX alone, 1 ethanol pretreatment alone, 2 ethanol pretreatment and DPCPX).

No differences in baseline systemic hemodynamics were observed between experimental groups (Table 1).

Twice daily consumption of ethanol for 12 weeks did not affect baseline systemic hemodynamics. Intravenous administration of drug vehicle or DPCPX in the presence or absence of ethanol pretreatment was also devoid of hemodynamic effects. Significant ($P < 0.05$) increases in LV end-diastolic pressure and

decreases in mean arterial and LV systolic pressures and $LV + dP/dt_{max}$ were observed during LAD occlusion and reperfusion. Heart rate was unchanged during LAD occlusion and reperfusion. No differences in systemic hemodynamics were observed between groups during occlusion or reperfusion.

The AAR was similar between groups (vehicle 40 ± 1 ; DPCPX 42 ± 3 ; ethanol pretreatment and vehicle 40 ± 2 ; ethanol pretreatment and DPCPX $42 \pm 2\%$ of the LV). Twelve week pretreatment with ethanol significantly reduced myocardial infarct size to $13 \pm 2\%$ of the AAR (Figs. 1 and 2) as compared to control experiments ($26 \pm 2\%$).

DPCPX abolished the protective effects of ethanol pretreatment ($30 \pm 3\%$) but had no effect in dogs that did not receive ethanol ($25 \pm 2\%$). Chronic intermittent ethanol ingestion for 12 weeks did not affect

Table 1
Systemic hemodynamics

	Baseline	Vehicle or OPCPX	Coronary occlusion	Reperfusion (min)		
				60	120	180
HR (min^{-1})						
Vehicle	128 \pm 4	128 \pm 4	117 \pm 7	117 \pm 6	120 \pm 4	122 \pm 5
DPCPX	129 \pm 7	126 \pm 7	127 \pm 5	126 \pm 3	126 \pm 5	121 \pm 6
Ethanol + vehicle	127 \pm 4	127 \pm 4	121 \pm 4	117 \pm 4	118 \pm 4	119 \pm 6
Ethanol + DPCPX	134 \pm 4	132 \pm 5	133 \pm 5	123 \pm 4	126 \pm 4	127 \pm 4
MAP (mmHg)						
Vehicle	104 \pm 3	104 \pm 3	88 \pm 3*	95 \pm 7	98 \pm 5	97 \pm 4
DPCPX	109 \pm 3	110 \pm 3	89 \pm 3*	90 \pm 2*	89 \pm 3*	91 \pm 2*
Ethanol + vehicle	111 \pm 6	111 \pm 6	97 \pm 7	96 \pm 6*	100 \pm 5	101 \pm 5
Ethanol + DPCPX	107 \pm 3	112 \pm 4	90 \pm 5	85 \pm 5*	92 \pm 6	94 \pm 4
LVSP (mmHg)						
Vehicle	117 \pm 3	117 \pm 3	93 \pm 4*	100 \pm 7	104 \pm 6	102 \pm 5
DPCPX	120 \pm 3	122 \pm 4	96 \pm 3*	93 \pm 3*	90 \pm 4*	96 \pm 2*
Ethanol + vehicle	122 \pm 7	122 \pm 7	105 \pm 6*	99 \pm 5*	104 \pm 4*	106 \pm 4*
Ethanol + DPCPX	118 \pm 4	125 \pm 4	84 \pm 6	87 \pm 6*	95 \pm 7*	98 \pm 5
LVEDP (mmHg)						
Vehicle	6 \pm 1	6 \pm 1	11 \pm 2*	15 \pm 2*	14 \pm 2*	15 \pm 1*
DPCPX	5 \pm 1	6 \pm 1	15 \pm 3*	13 \pm 2*	11 \pm 3	12 \pm 3
Ethanol + vehicle	7 \pm 1	7 \pm 1	11 \pm 2*	8 \pm 2	9 \pm 2	8 \pm 2
Ethanol + DPCPX	5 \pm 1	6 \pm 1	15 \pm 2*	17 \pm 2*	18 \pm 2*	14 \pm 2*
+dP/dt _{max} (mmHg·s ⁻¹)						
Vehicle	1892 \pm 159	1892 \pm 159	1561 \pm 53	1445 \pm 70	1504 \pm 108	1279 \pm 55*
DPCPX	1967 \pm 120	1927 \pm 114	1530 \pm 72*	1576 \pm 51*	1555 \pm 59*	1456 \pm 70*
Ethanol + vehicle	1921 \pm 172	1921 \pm 172	1484 \pm 172*	1366 \pm 83*	1337 \pm 102*	1435 \pm 84*
Ethanol + DPCPX	1922 \pm 111	1915 \pm 118	1663 \pm 191	1362 \pm 107*	1470 \pm 103*	1480 \pm 99*

Data are mean \pm SEM; $n = 7$ in each group.

*Significantly ($P < 0.05$) different from vehicle or DPCPX.

Abbreviations: HR = heart rate; MAP = mean arterial pressure; LVSP and LVEDP = left ventricular systolic and end-diastolic pressures, respectively; +dP/dt_{max} = peak rate of increase of left ventricular pressure; DPCPX = 8-cyclopentyl-1,3-dipropylxanthine.




Fig. 1. Myocardial infarct size (IF) expressed as a percentage of the area of the left ventricle at risk (AAR). Abbreviation: DPCPX=8-cyclopentyl-1,3-dipropylxanthine. *Significantly ($P<0.05$) different from VEHICLE; †Significantly ($P<0.05$) different from DPCPX; §Significantly ($P<0.05$) different from ETHANOL+DPCPX.

baseline regional myocardial perfusion. No differences in transmural coronary collateral blood flow (Table 2) were observed between groups.

4. Discussion

Previous investigations from this [3,4] and other laboratories [5,10–12,17] have demonstrated that chronic, moderate ethanol consumption protects myocardium from ischemic injury in isolated and intact hearts. The results of these experiments obtained in several animal species are supported by recent clinical observations that chronic ingestion of small amounts of ethanol enhances survival after myocardial infarction in humans [1,2]. The precise mechanisms responsible for this beneficial effect remain to be completely defined, but a signal transduction cascade has been implicated in CEPC that is remarkably similar to a major pathway responsible for IPC. The present results confirm the previous findings indicating that chronic, intermittent ethanol ingestion reduces the extent of myocardial infarction associated with prolonged coronary artery occlusion and reperfusion. The reduction in infarct size produced by chronic ethanol ingestion observed in dogs in the present and previous studies [3,4] occurred independent of alterations in systemic hemodynamics and coronary collateral blood flow. Other studies conducted in rats [5,12,17] and guinea pigs [10–12] did not quantify transmural myocardial perfusion




Table 2
Transmural myocardial blood flow in the ischemic and normal regions (ml/min/g)

	Vehicle of DPCPX	Coronary occlusion	Reperfusion
Ischemic region			
Vehicle	1.08±0.11	0.08±0.01*	1.22±0.14
DPCPX	1.25±0.20	0.04±0.01*	1.58±0.31
Ethanol + vehicle	0.88±0.15	0.05±0.01*	0.90±0.11
Ethanol + DPCPX	0.80±0.05	0.06±0.01*	1.00±0.15
Normal region			
Vehicle	1.22±0.11	1.12±0.11	1.23±0.14
DPCPX	1.55±0.27	1.21±0.24	1.47±0.30
Ethanol + vehicle	1.34±0.17	0.98±0.14	0.94±0.11
Ethanol + DPCPX	1.05±0.16	0.93±0.11	1.18±0.14

Data are mean±SEM; n=7 in each group.

*Significantly ($P<0.05$) different from vehicle or DPCPX.

Abbreviation: DPCPX = 8-cyclopentyl-1,3-dipropylxanthine.

demonstrated a role for adenosine receptors in chronic ethanol-induced protection against myocardial ischemic injury. Isolated hearts from guinea pigs exposed to low concentrations (2.5 or 5%) of ethanol in their drinking water for 12 weeks before experimentation demonstrated improved functional recovery (peak developed LV pressure) and decreased myocardial necrosis (creatinine kinase release) after 45 min of no-flow ischemia as compared with controls [10,12]. These salutary effects were inhibited by the nonselective adenosine receptor antagonist 8-(*p*-sulphophenyl)theophylline (SPT) and DPCPX but not by the selective A_2 antagonist 3,7-dimethyl-1-propargylxanthine (DMPX) [10,12]. The results with a 12 week exposure to ethanol were similar to those obtained with IPC produced by 2 min of global ischemia followed by 5 min of reperfusion before no-flow ischemia in the absence of ethanol pretreatment. The present and previous [10,12] results are also partially supported by recent findings demonstrating that adenosine-induced myocardial protection against excessive catecholamine stimulation may be enhanced by chronic ethanol consumption in rats [18]. Hearts chronically exposed to ethanol demonstrated increased sensitivity to the protective effects of adenosine despite reductions in total myocardial adenosine release measured in coronary venous blood. These findings suggested that chronic ethanol ingestion may produce an upregulation or enhanced responsiveness of myocardial adenosine receptors

[18]. Such phenomena may be associated with increased myocardial protection against ischemic injury because of the well-established link between A_1 receptors and K_{ATP} channels [7,19]. This intriguing hypothesis will require additional research to confirm, however. Extrapolation of the conclusions of Fenton and Chung [18] to the present results should also be qualified because larger amounts of ethanol (36% of total daily caloric intake) were chronically ingested for a more prolonged duration (8 months) by rats in this previous study. Whether administration of DPCPX during the preconditioning period prevents the development of CEPC also remains unknown.

In contrast to the continuous administration of ethanol in drinking water described in many previous studies [5,10–12,17], the present investigation used intermittent administration of ethanol with ad libitum access to fresh drinking water during the pretreatment period. Myocardial protection associated with this method of administration occurred after ethanol had been withheld the night before experimentation, in contrast to the presence of an average blood ethanol concentration of 3 mM immediately before experimentation previously described in one study [5]. Acute exposure to ethanol concentrations as low as 10 mM has been shown to produce direct protective effects against ischemic damage in ventricular myocytes and isolated hearts [20,21] and enhance the functional recovery of stunned myocardium in vivo [22]. There is also experimental evidence in neural tissue indicating that acute administration of ethanol may directly activate A_1 receptors [23] by increasing the extracellular concentration of adenosine [24]. Thus, it remains possible that previous results implicating the A_1 receptor in CEPC obtained in small mammals continuously exposed to ethanol may have been influenced to some degree by the presence of ethanol immediately before experimentation. However, acute administration of intoxicating doses of ethanol immediately before coronary artery occlusion did not reduce infarct size in similar canine model [25]. These findings suggest that chronic ethanol ingestion may be required for myocardial protection to occur.

The present results should be interpreted within the constraints of several potential limitations. The specificity of DPCPX for the A_1 receptor has not been definitively established in canine myocardium, but

this compound has been shown to be 700 times more selective for A_1 versus A_2 receptors using radioligand binding studies and in vitro functional assays [26]. Previous studies also indicate that the dose of DPCPX (0.8 mg/kg) used in the present investigation attenuates isoflurane-induced improvements in the functional recovery of stunned myocardium [15] and abolishes IPC in the canine heart [19] via A_1 receptor blockade. It is unlikely that myocardial protection produced by chronic, intermittent ethanol ingestion and the abolition of this beneficial effect by DPCPX were related to differential alterations in myocardial oxygen consumption because systemic hemodynamics were very similar between experimental groups. Nevertheless, coronary sinus oxygen tension was not specifically measured and myocardial oxygen consumption was not directly calculated in the present investigation. Whether DPCPX acutely alters the well-known beneficial effects of chronic, intermittent ethanol ingestion on platelet function and lipid metabolism also cannot be completely excluded as a potential explanation for observed findings. The present results also require qualification because oral ingestion of the dose of ethanol used in this investigation may produce greater blood ethanol concentrations in humans than dogs, although the pharmacokinetics of ethanol appear to be relatively similar between these species.

In summary, the present results demonstrate that chronic, intermittent consumption of small amounts of ethanol reduce myocardial infarct size in dogs independent of alterations in hemodynamics or transmural myocardial perfusion. This protective effect was abolished by the selective A_1 antagonist DPCPX, indicating that the A_1 receptor mediates CEPC in vivo.

Acknowledgements

This work was supported in part by grants AA-12331 (Dr Pagel), HL-03690 (Dr Kersten), HL-63705 (Dr Kersten), HL-54820 (Dr Warltier), and GM-08377 (Dr Warltier) from the United States Public Health Service, Bethesda, Maryland. The authors thank Mr David A. Schwabe and Mr John P. Tessmer for technical assistance.

References

- [1] Wannamethee G, Whincup PH, Shaper AG, Walker M, MacFarlane PW. Factors determining case fatality in myocardial infarction 'Who dies in a heart attack?'. *Br Heart J* 1995;74:324–31.
- [2] Mukamal KJ, Maclure M, Muller JE, Sherwood JB, Mittleman MA. Prior alcohol consumption and mortality following acute myocardial infarction. *J Am Med Assoc* 2001;285:1965–70.
- [3] Pagel PS, Toller WG, Gross ER, Gare M, Kersten JR, Warltier DC. K_{ATP} channels mediate the beneficial effects of chronic ethanol ingestion. *Am J Physiol Heart Circ Physiol* 2000;279:H2574–9.
- [4] Pagel PS, Krolikowski JG, Kehl F, Mraovic B, Kersten JR, Warltier DC. The role of mitochondrial and sarcolemmal K_{ATP} channels in canine ethanol-induced preconditioning in vivo. *Anesth Analg* 2002;94:841–8.
- [5] Zhu P, Zhou HZ, Gray MO. Chronic ethanol-induced myocardial protection requires activation of mitochondrial KATP channels. *J Mol Cell Cardiol* 2000;32:2091–5.
- [6] Liu Y, Sato T, O'Rourke B, Marban E. Mitochondrial ATP-dependent potassium channels. Novel effectors of cardioprotection? *Circulation* 1998;97:2463–9.
- [7] Sato T, Sasaki N, O'Rourke B, Marban E. Adenosine primes the opening of mitochondrial ATP-sensitive potassium channels. A key step in ischemic preconditioning? *Circulation* 2000;102:800–5.
- [8] Schultz JE, Hsu AK, Barbieri JT, Li PL, Gross GJ. Pertussis toxin abolishes the cardioprotective effect of ischemic preconditioning in the intact rat heart. *Am J Physiol Heart Circ Physiol* 1998;44:H495–500.
- [9] Ping P, Zhang J, Qiu Y, Tang XL, Manchikalapudi S, Cao X, Bolli R. Ischemic preconditioning induces selective translocation of protein kinase C isoforms ϵ and η in the heart of conscious rabbits without subcellular redistribution of total protein kinase C activity. *Circ Res* 1997;81:404–14.
- [10] Miyamae M, Diamond I, Weiner MW, Camacho SA, Figueredo VM. Regular alcohol consumption mimics cardiac preconditioning by protecting against ischemia-reperfusion injury. *Proc Natl Acad Sci USA* 1997;94:3235–9.
- [11] Miyamae M, Rodriguez MM, Camacho SA, Diamond I, Mochly-Rosen D, Figueredo VM. Activation of ϵ protein kinase C correlates with a cardioprotective effect of regular ethanol consumption. *Proc Natl Acad Sci USA* 1998;95:8262–7.
- [12] Miyamae M, Camacho SA, Zhou H-Z, Diamond I, Figueredo VM. Alcohol consumption reduces ischemia-reperfusion injury by species-specific signaling in guinea pigs and rats. *Am J Physiol Heart Circ Physiol* 1998;44:H50–6.
- [13] Ferguson ER, Blachley JD, Carter NW, Knochel JP. Derangements of muscle composition, ion transport, and oxygen consumption in chronically alcoholic dogs. *Am J Physiol* 1984;246:F700–9.
- [14] Kersten JR, Schmeling TJ, Pagel PS, Gross GJ, Warltier DC. Isoflurane mimics ischemic preconditioning via activation of K_{ATP} channels: reduction of myocardial infarct size with an acute memory phase. *Anesthesiology* 1997;87:361–70.
- [15] Kersten JR, Orth KG, Pagel PS, Mei DA, Gross GJ, Warltier DC. Role of adenosine in isoflurane-induced cardioprotection. *Anesthesiology* 1997;86:1128–39.
- [16] Gross GJ, Auchampach JA. Blockade of ATP-sensitive potassium channels prevents myocardial preconditioning in dogs. *Circ Res* 1992;70:223–33.
- [17] McDonough KH. Chronic alcohol consumption causes accelerated myocardial preconditioning to ischemia-reperfusion injury. *Alcohol Clin Exp Res* 1997;21:869–73.

- [18] Fenton RA, Chung ES. Chronic ethanol enhances adenosine antiadrenergic actions in the isolated rat heart. *Alcohol Clin Exp Res* 2001;25:968–75.
- [19] Auchampach JA, Gross GJ. Adenosine A₁ receptors, K_{ATP} channels and ischemic preconditioning in dogs. *Am J Physiol Heart Circ Physiol* 1993;33:H1327–36.
- [20] Chen CH, Gray MO, Mochly-Rosen D. Cardioprotection from ischemia by a brief exposure to physiological levels of ethanol: role of epsilon protein kinase C. *Proc Natl Acad Sci USA* 1999;96:12784–9.
- [21] Krenz M, Baines CP, Yang XM, Heusch G, Cohen MV, Downey JM. Acute ethanol exposure fails to elicit preconditioning-like protection in in situ rabbit hearts because of its continued presence during ischemia. *J Am Coll Cardiol* 2001;37:601–7.
- [22] Gross ER, Gare M, Toller WG, Kersten JR, Wartier DC, Pagel PS. Ethanol enhances the functional recovery of stunned myocardium independent of K_{ATP} channels in dogs. *Anesth Analg* 2001;92:299–305.
- [23] Diamond I, Gordon AS. Cellular and molecular neuroscience of alcoholism. *Physiol Rev* 1997;77:1–20.
- [24] Nagy LE, Diamond I, Casso DJ, Franklin C, Gordon AS. Ethanol increases extracellular adenosine by inhibiting adenosine uptake via the nucleoside transporter. *J Biol Chem* 1990;265:1946–51.
- [25] Itoya M, Morrison JD, Downey HF. Effect of ethanol on myocardial infarct size in a canine model of coronary artery occlusion-reperfusion. *Mol Cell Biochem* 1998;186:35–41.
- [26] Lohse MJ, Klotz KN, Lindenborn-Fotinos J, Reddington M, Schwabe U, Olsson RA. 8-Cyclopentyl-1,3-dipropylxanthine (DPCPS)—a selective high affinity antagonist radioligand for A₁ adenosine receptors. *Naunyn Schmiedebergs Arch Pharmacol* 1987;336:204–10.